## For Research Use Only UBXN2B Polyclonal antibody

Catalog Number:25141-1-AP




| Basic Information  | Catalog Number:<br>25141-1-AP  | GenBank Accession Number:<br>BC113645                     | Purification Method:<br>Antigen affinity purification                                 |                 |
|--|--|---|---|-----------------|
|  | Size:<br>1000 ug/ml<br>Source:<br>Rabbit   | GeneID (NCBI):<br>137886<br>UNIPROT ID:<br>Q14CS0         | Recommended Dilutions:<br>WB 1:2000-1:10000<br>IHC 1:400-1:1600<br>IF/ICC 1:200-1:800 |                 |
|  |  |   |   | lsotype:<br>IgG |
|  | Immunogen Catalog Number:<br>AG18259   | Calculated MW:<br>331 aa, 37 kDa                          |   |                 |
|  |  | Observed MW:<br>37 kDa                                    |   |                 |
|  | Applications   | Tested Applications: Positive C<br>WB, IHC, IF/ICC, ELISA |   |                 |
| Species Specificity:<br>human, mouse   |  |   | WB : Neuro-2a cells,<br>IHC : human ovary cancer tissue,                              |                 |
| Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0 |  | Neuro-2a cells,   |   |                 |
| Background Information   | UBXN2B, also called p37, is a cofactor of the p97 AAA ATPase. UBXN2B regulates spindle orientation during mitosis<br>by limiting the levels of cortical NuMA (PMID: 29222185). |   |   |                 |
| Storage  | Storage:Store at -20°C. Stable for one year after shipment.Storage Buffer:PBS with 0.02% sodium azide and 50% glycerol pH 7.3.Aliquoting is unnecessary for -20°C storage      |   |   |                 |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**



Neuro-2a cells lysates were subjected to SDS PAGE followed by western blot with 25141-1-AP (UBXN28 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 25141-1-AP (UBXN2B antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using UBXN2B antibody (25141-1-AP) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).