For Research Use Only

NAT12 Polyclonal antibody

Catalog Number: 25149-1-AP



Purification Method:

WB 1:1000-1:6000

Antigen affinity purification

Recommended Dilutions:

Basic Information

 Catalog Number:
 GenBank Accession Number:

 25149-1-AP
 BC118589

 Size:
 GeneID (NCBI):

 160 μg/ml
 122830

 Source:
 UNIPROT ID:

 Rabbit
 Q147X3

UNIPROT ID: IF/ICC 1:200-1:800 Q147X3
Full Name:

Isotype: IgG

IgG N-acetyltransferase 12 (GCN5-related,

Immunogen Catalog Number: putative)

AG18435 Calculated MW:

362 aa, 39 kDa Observed MW: 40 kDa

Applications

Tested Applications:

IF/ICC, WB, ELISA

Species Specificity:
human

Positive Controls:

WB: HEK-293 cells,

IF/ICC: U2OS cells,

Background Information

N-terminal acetylation (Nt-acetylation) by N-terminal acetyltransferases (NATs) is one of the most common protein modifications in eukaryotes. The NatC complex represents one of three major NATs of which the substrate profile remains largely unexplored. NAT12/NAA30 (N-alpha-acetyltransferase 30) is the catalytic subunit of the NAT-C complex (PMID: 27694331, 26292663). The expected sizes of the two isoforms of the protein were 35 and 39 kDa. Western blots detected two bands of NAA12, the expected ~40 kDa band and a possibly dimer at ~70 kDa.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25149-1-AP (NAT12 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using NAT12 antibody (25149-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).