For Research Use Only

SFMBT2 Polyclonal antibody

Catalog Number:25256-1-AP

3 Publications



Basic Information

Catalog Number: 25256-1-AP

Size: $600 \mu g/ml$ Source: Rabbit Isotype:

Scm-like with four mbt domains 2 Calculated MW:

Immunogen Catalog Number:

AG19379

894 aa, 101 kDa Observed MW: 100 kDa

BC152430

57713

GeneID (NCBI):

UNIPROT ID:

Q5VUG0 Full Name:

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:3000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, CoIP

Species Specificity: mouse, rat **Cited Species:** human, mouse

Positive Controls:

WB: mouse brain tissue, NIH/3T3 cells, mouse spleen tissue, rat brain tissue

Background Information

SFMBT2 (Scm like with four mbt domains 2), also named as KIAA1617. It is located in Nucleus and ubiquitous expression in thyroid and ovary. The calculated molecular weight of SFMBT2 is 100 kDa. Although mammalian SFMBTs have similar structural features with PcG protein dSFMBT, biological function and target genes of SFMBT have been not investigated well. Lee K and Na W investigated biological function of human SFMBT2 that interacted with YY1 using AR (androgen receptor)-negative DU145 prostate cancer cells. They found that human SFMBT2 represses HOXB13 gene expression via its association with methylated histones H3 and H4 that are transcriptional repression marks. Furthermore, their findings indicate that SFMBT2 may be involved in regulation of cell growth of DU145 prostate cancer cells (PMID: 23385818).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|----------------|-------------|
| Safdar Hussain | 30272322 | Int J Mol Med | |
| Le-Yun Wang | 32559418 | Cell Stem Cell | |
| Qingpeng Xie | 38734627 | Biol Direct | WB,CoIP |

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25256-1-AP (SFMBT2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.