For Research Use Only

ORP1L Polyclonal antibody

Catalog Number: 25268-1-AP



Basic Information

Catalog Number:

25268-1-AP

Size: 350 ug/ml Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

NM_018030 GeneID (NCBI): 114876 UNIPROT ID:

Q9BXW6 Full Name:

охysterol binding protein-like 1А

Calculated MW: 108 kDa Observed MW: 50 kDa, 105-110 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

Applications

Tested Applications:

WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: mouse brain tissue, rat brain tissue

Background Information

ORP1L (Oxysterol-binding protein-related protein 1L) is a lipid transfer protein that belongs to the family of OSBP-related proteins (ORPs). It is recruited to lysosomes via its ankyrin repeat domains that bind to Rab7(PMID: 34504082). ORP1L also possesses an FFAT motif, which allows it to interact with the ER resident VAP proteins to aid in establishing the ER-lysosomes contacts. ORP1L is expressed in various tissues, with particularly high expression in the brain, lungs, and macrophages. ORP1L plays a crucial role in cholesterol homeostasis. Disruption of ORP1L function can lead to a block in cholesterol transport to the ER, resulting in the accumulation of cholesterol-enriched LELs in the Golgi/perinuclear region. This can disrupt cellular cholesterol balance, potentially contributing to various metabolic diseases. This suggests that ORP1L is necessary for the normal functioning of cholesterol export from the endo-lysosomal system to the ER, and its dysfunction could be implicated in the progression of diseases related to cholesterol metabolism dysregulation(PMID: 28564600). The molecular weight of ORP1L is 50 kDa.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25268-1-AP (ORP1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.