

CHD3 Polyclonal antibody

Catalog Number: 25335-1-AP

1 Publications

Basic Information

Catalog Number:

25335-1-AP

Size:

900 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_005852

GeneID (NCBI):

1107

UNIPROT ID:

Q12873

Full Name:

chromodomain helicase DNA binding protein 3

Calculated MW:

227 kDa

Observed MW:

240 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : MCF-7 cells, HEK-293 cells, mouse brain tissue

IHC : mouse brain tissue, human colon tissue

Background Information

CHD3, also named as Mi2 ALPHA or Zinc finger helicase, is a 2000 amino acid protein, which contains one helicase C-terminal domain, one helicase ATP-binding domain, and two chromo domain. CHD3 localizes in the nucleus and cytoplasm. CHD3 is widely expressed. CHD3 is a component of the histone deacetylase NuRD complex, which participates in the remodeling of chromatin by deacetylating histones. CHD3 is required for anchoring centrosomal pericentric in both interphase and mitosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Xue Chen	38676947	Nucleic Acids Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 25335-1-AP (CHD3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 25335-1-AP (CHD3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 25335-1-AP (CHD3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 25335-1-AP (CHD3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 25335-1-AP (CHD3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 25335-1-AP (CHD3 Antibody) at dilution of 1:50 (under 40x lens).