For Research Use Only

NOX5 Polyclonal antibody Catalog Number:25350-1-AP 12 Publications




Basic Information	Catalog Number: 25350-1-AP			Purification Meth		
	Concentration:	GenelD (NCBI):		Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000		
	400 ug/ml	79400				
	Source: Rabbit	UNIPROT ID: Q96PH1		IHC 1:50-1:500		
	Isotype:	Full Name:	-			
	IgG Immunogen Catalog Number: AG18673		NADPH oxidase, EF-hand calcium binding domain 5			
		Calculated MW: 765 aa, 86 kDa				
		Observed MW: 82-86 kDa				
Applications	Tested Applications:	Positive Controls:				
	WB, IHC, ELISA	WB: Jurkat Cetts, Kaji Cetts				
	Cited Applications: WB, CoIP, PLA		IHC : human spleen tissue,			
	Species Specificity: human					
	Cited Species: human, pig, monkey					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	NOX5 (NADPH oxidase, EF-hand functions as an H+ channel in a C activation and human aortic smo is implicated in human spermate NOX5- α , - β , - γ , - δ , - ε and - ζ Uniprot).	Ca(2+)-dependent man both muscle cell (HAS bozoa motility. This pro	nner. It plays an impo MC) proliferation. NO otein has 6 isoforms p	ortant role in PDGF- 0X5 is the main sou produced by alterna	induced JAK/STAT rce of superoxide a ntive splicing, nam	
Notable Publications	Author	Pubmed ID 3	Journal		Application	
	Jian Wu	31727371 l	Biochem Biophys Res	Commun	WB	
	Elizabeth A Sweeny		Free Radic Biol Med		WB	
			Biomed Pharmacothe	r	WB,PLA,CoIP	
	Qi Liu	32738790 I	biomed mannacoure			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 25350-1-AP (NOX5 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 25350-1-AP (NOX5 Antibody) at dilution of 1:200 (under 10x lens).