For Research Use Only

TTC35 Polyclonal antibody

Catalog Number: 25443-1-AP

Featured Product

13 Publications

BC021667

9694

GeneID (NCBI):

UNIPROT ID:

Q15006 Full Name:

GenBank Accession Number:

tetratricopeptide repeat domain 35



Basic Information

Catalog Number: 25443-1-AP

Size: 1200 µg/ml Source:

Rabbit Isotype:

AG22196

Immunogen Catalog Number:

297 aa, 35 kDa Observed MW:

Calculated MW:

35 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications: IP, WB, ELISA

Cited Applications: WB, IP, IF

Species Specificity: human, mouse, rat **Cited Species:**

human, mouse

Positive Controls:

WB: Jurkat cells, mouse heart tissue, K-562 cells, MCF-7 cells, NIH/3T3 cells, mouse spleen tissue, human placenta tissue, PC-3 cells, SKOV-3 cells

IP: mouse heart tissue,

Background Information

TPR repeat protein 35 (TTC35), also named as tetratricopeptide repeat domain 35, is a 297 amino acid protein, which contains three TPR repeats and belongs to the EMC2 family. TTC35 localizes in the nucleus and cytoplasm. TTC35 is a component of the ER membrane protein complex and may be a putative O-linked glycosyl transferase.

Notable Publications

Author	Pubmed ID	Journal	Application
Ashley M Ngo	31516121	Elife	WB
Patrick J Chitwood	30415835	Cell	WB
Cary R Boyd-Shiwarski	36318922	Cell	IF

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 25443-1-AP (TTC35 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-TTC35 (IP:25443-1-AP, 3ug; Detection:25443-1-AP 1:600) with mouse heart tissue lysate 4000ug.



Various lysates were subjected to SDS PAGE followed by western blot with 25443-1-AP (TTC35 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.