

For Research Use Only

P27; KIP1 Polyclonal antibody, PBS Only

Catalog Number: 25614-1-PBS

Featured Product



Basic Information

Catalog Number:

25614-1-PBS

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG22582

GenBank Accession Number:

BC001971

GeneID (NCBI):

1027

UNIPROT ID:

P46527

Full Name:

cyclin-dependent kinase inhibitor 1B
(p27, Kip1)

Calculated MW:

198 aa, 22 kDa

Observed MW:

27 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, FC (Intra), IP, Indirect ELISA

Species Specificity:

human, mouse

Background Information

CDKN1B, also named as P27 or KIP1, is a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. P27 binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controlling cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Downregulation of P27 has been implicated in the progression of several malignancies, including lung cancer, hepatocellular carcinoma, salivary cancer, oral squamous cell carcinomas, and gastric cancer.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 25614-1-AP (P27; KIP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 25614-1-AP (P27; KIP1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 25614-1-AP (P27; KIP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.




WB result of P27; KIP1 antibody (25614-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-P27; KIP1 transfected HeLa cells. This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 25614-1-AP (P27; KIP1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 25614-1-AP (P27; KIP1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.





Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using P27; KIP1 antibody (25614-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using P27; KIP1 antibody (25614-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human P27; KIP1 (25614-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 25614-1-PBS in a different storage buffer formulation.



Normal Rabbit IgG (020000-01)
P27; KIP1 Antibody (02014-1-AP)

IP result of anti-P27; KIP1 (IP:25614-1-AP, 4ug;
Detection:25614-1-AP 1:1000) with NIH/3T3 cells
lysate 1160 ug. This data was developed using the
same antibody clone with 25614-1-PBS in a
different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed
HepG2 cells using P27; KIP1 antibody (25614-1-AP)
at dilution of 1:200 and CoraLite®594-Conjugated
Goat Anti-Rabbit IgG(H+L) (SA00013-4), CL488-
Phalloidin (green). This data was developed using
the same antibody clone with 25614-1-PBS in a
different storage buffer formulation.