For Research Use Only

CHCHD10 Polyclonal antibody

Catalog Number:25671-1-AP

Featured Product

20 Publications




Basic Information	Catalog Number: 25671-1-AP			Purification Method: Antigen affinity purification				
	Concentration: 600 ug/ml	GenelD (NCB 400916	l):	Recommended Dilutions: WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total				
	Source:	UNIPROT ID:						
	Rabbit Isotype: IgG Immunogen Catalog Number: AG22598	Q8WYQ3 Full Name: coiled-coil-helix-coiled-coil-helix domain containing 10 Calculated MW: 14 kDa		protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800				
					Observed MV 14 kDa	t.		
					Applications	Tested Applications: WB, IHC, IF/ICC, IP, CoIP, ELISA	Positive Contr	
		Cited Applications: www.inc.29 WB; IHC, IF, IP IP : mouse h Species Specificity: IHC : humar human, mouse, rat IHC : humar		5 cells, mouse brain tissue, HAP1, mouse rat heart tissue				
eart tissue, HAP1 cells								
i kidney tissue, human heart tissue								
Cited Species: IF/ICC : HeLa human, mouse, chicken			o cells, HAP1					
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0								
Background Information	involved in mitochondrial genon	ne stability and ma	aintenance of cristae ju	the intermembrane space and is likely nctions. Mutations in CHCHD10 have orbid with amyotrophic lateral sclerosis				
Notable Publications	Author	Pubmed ID	Journal	Application				
	Meng Ding	35709007	Diabetes	WB				
	Wei Zhou	30496485	Hum Mol Genet	WB,IF				
	Mai K Nguyen	34791217	Hum Mol Genet	WB,IF				
	Storage:							


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Co-IP result of anti-CHCHD10 (IP:25671-1-AP, 4ug; Detection:19424-1-AP 1:600) with HepG2 cells lysate 1400ug.




Various lysates were subjected to SDS PAGE followed by western blot with 25671-1-AP (CHCHD10 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




HEK-293 cells were subjected to SDS PAGE followed by western blot with 25671-1-AP (CHCHD10 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




HAP1 (WT and CHCHD10 KO) lysates prepared with RIPA buffer, 50 μ g protein loaded. 25671-1-AP incubated at 1:1000 at 4°C overnight in 5% milk in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




WB result of CHCHD10 antibody (25671-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CHCHD10 transfected HEK-293 cells.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25671-1-AP (CHCHD10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25671-1-AP (CHCHD10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CHCHD10 antibody (25671-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




HAP1 WT cells (yellow outline) and CHCHD10 KO cells (red outline) labelled with a green or a far-red fluorescence dye, respectively. Cells fixed with 4% PFA and stained with 25671-1-AP at 1:300 plus DAPI. Bars = 10 μ m. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.



SM=10% starting material; UB=10% unbound fraction; IP=immunoprecipitate





HAP1 lysates prepared and IP of CHCHD10 performed using 1.0 μ g of 25671-1-AP coupled to protein A- Sepharose beads. The Ponceau stained transfers of each blot are shown. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.

IP result of anti-CHCHD10 (IP:25671-1-AP, 4ug; Detection:25671-1-AP 1:600) with mouse heart tissue lysate 4000ug. Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CHCHD10 antibody (25671-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).