

For Research Use Only

CD63 Polyclonal antibody

Catalog Number: 25682-1-AP

Featured Product

481 Publications



Basic Information

Catalog Number:

25682-1-AP

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG19690

GenBank Accession Number:

BC002349

GeneID (NCBI):

967

UNIPROT ID:

P08962

Full Name:

CD63 molecule

Calculated MW:

26 kDa

Observed MW:

30-60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:40000

IHC: 1:400-1:1600

IF-P: 1:50-1:500

IF/ICC: 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications:

WB, IHC, IF, IP, ChIP

Species Specificity:

human

Cited Species:

human, pig, rabbit, monkey, chicken, zebrafish, hamster, goat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human urine exosomes tissue, A375 cells, human serum exosomes, THP-1 cells, human peripheral blood platelets

IHC: human malignant melanoma tissue, human lymphoma tissue, human tonsillitis tissue

IF-P: human malignant melanoma tissue,

IF/ICC: A431 cells,

Background Information

CD63 is a 30-60 kDa lysosomal membrane protein that belongs to the tetraspanin family. This protein plays many important roles in immuno-physiological functions. It mediates signal transduction events that play a role in the regulation of cell development, activation, growth, and motility. CD63 is expressed on activated platelets, thus it may function as a blood platelet activation marker. CD63 is a lysosomal membrane glycoprotein that is translocated to plasma membrane after platelet activation. The CD63 tetraspanin is highly expressed in the early stages of melanoma and decreases in advanced lesions, suggesting it as a possible suppressor of tumor progression. Deficiency of this protein is associated with Hermansky-Pudlak syndrome.

Notable Publications

Author	Pubmed ID	Journal	Application
Yong Chen	36224027	Front Biosci (Landmark Ed)	WB
Xinyan Zhou	36171212	Nat Commun	WB
Jingming Zhuang	36161709	Andrology	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human urine exosomes tissue were subjected to SDS PAGE followed by western blot with 25682-1-AP (CD63 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human malignant melanoma tissue slide using 25682-1-AP (CD63 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human malignant melanoma tissue slide using 25682-1-AP (CD63 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human malignant melanoma tissue using CD63 antibody (25682-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed human malignant melanoma tissue using CD63 antibody (25682-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 25682-1-AP (CD63 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 25682-1-AP (CD63 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using CD63 antibody (25682-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).