### For Research Use Only

# Chk1 Polyclonal antibody

Catalog Number: 25887-1-AP

**Featured Product** 

22 Publications



**Basic Information** 

Catalog Number: 25887-1-AP

BC004202 GeneID (NCBI): 1111

Calculated MW:

Full Name: IP 0.5-4.0 ug for 1.0-3.0 mg of total CHK1 checkpoint homolog (S. pombe) protein lysate IHC 1:100-1:400

GenBank Accession Number:

IgG 54 kDa
Immunogen Catalog Number: Observed MW:
AG22993 55 kDa

Applications

**Tested Applications:** 

Size: 450 μg/ml

Source:

Rabbit

Isotype:

FC, IF, IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, IP, WB Species Specificity:

Cited Species: human, mouse

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293T cells, HeLa cells, K-562 cells, MCF-7

**Purification Method:** 

WB 1:1000-1:6000

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cells

IP: HEK-293T cells,
IHC: human kidney tissue,
IF: HEK-293T cells,

## **Background Information**

CHEK1(Checkpoint kinase-1) is also named as CHK1 and belongs to the protein kinase superfamily. It is implicated in a circuit in which it activates checkpoints, DNA repair and proliferating cell nuclear antigen and FANCD2 monoubiquitinylation(PMID:21389083). CHEK1 protects vertebrate cells against spontaneous chromosome missegregation and is required to sustain anaphase delay when spindle function is disrupted by taxol(PMID:17276342). It has 3 isoforms produced by alternative splicing with the molecular mass of 54 kDa, 44 kDa and 50 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yeunting Hsieh	32988875	Anticancer Res	WB
Jingyuan Sun	33087136	J Exp Clin Cancer Res	WB,IF
Tai-Hsin Tsai	34858180	Front Pharmacol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 25887-1-AP (Chk1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Methanol) fixed HEK-293T cells using Chk1 antibody (25887-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP Result of anti-CHK1 (IP:25887-1-AP, 4ug; Detection:25887-1-AP 1:600) with HEK-293T cells lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25887-1-AP (CHK1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25887-1-AP (CHK1 Antibody) at dilution of 1:200 (under 40x lens).



1X10^6 HEK-293T cells were intracellularly stained with 0.2 ug Anti-Human Chk1 (25887-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).