

For Research Use Only

# CCR7 Polyclonal antibody

Catalog Number: 25898-1-AP

19 Publications



## Basic Information

Catalog Number:

25898-1-AP

Concentration:

300 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG22941

GenBank Accession Number:

BC035343

GeneID (NCBI):

1236

UNIPROT ID:

P32248

Full Name:

chemokine (C-C motif) receptor 7

Calculated MW:

378 aa, 43 kDa

Observed MW:

40-43 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:1000

IHC: 1:50-1:500

## Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, rat

Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : MCF-7 cells, C6 cells, K-562 cells, T-47D cells

IHC : human tonsillitis tissue,

## Background Information

C-C chemokine receptor type 7 (CCR7), which belongs to the G-protein coupled receptor 1 family, is also a receptor for the MIP-3-beta chemokine, and probably acts as a mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions. CCR7 is specially expressed in various lymphoid tissues and activated B- and T-lymphocytes, and can be strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7.

## Notable Publications

Author	Pubmed ID	Journal	Application
Mingzhao Zhang	32943109	Stem Cell Res Ther	IF
Xiaolei Shi	36217946	J Agric Food Chem	WB
Zhengmin Liang	35603910	Virulence	IHC

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 25898-1-AP (CCR7 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



MCF-7 cells were subjected to SDS PAGE followed by western blot with 25898-1-AP (CCR7 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 25898-1-AP (CCR7 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 25898-1-AP (CCR7 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).