

For Research Use Only

# SNAT2 Polyclonal antibody

Catalog Number: 25928-1-AP



## Basic Information

<b>Catalog Number:</b> 25928-1-AP	<b>GenBank Accession Number:</b> BC040342	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 500 ug/ml	<b>GeneID (NCBI):</b> 54407	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:200-1:800
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q96QD8	
<b>Isotype:</b> IgG	<b>Full Name:</b> solute carrier family 38, member 2	
<b>Immunogen Catalog Number:</b> AG23082	<b>Observed MW:</b> 45 kDa, 56 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human, mouse, rat	<b>WB :</b> 37°C incubated mouse brain tissue,
<b>Note-IHC:</b> suggested antigen retrieval with <b>TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	<b>IHC :</b> mouse pancreas tissue, human cervical cancer tissue, human kidney tissue, rat pancreas tissue, rat kidney tissue

## Background Information

SNAT2 (Sodium-dependent neutral amino acid transporter-2), the ubiquitous member of SLC38 family, accounts for the activity of transport system A for neutral amino acids in most mammalian tissues (PMID: 16734764). SNAT2 mediates uptake of neutral  $\alpha$ -amino acids and are expressed in central neurons (PMID: 19240036). SNAT2 is expressed in breast cancer cell lines. SLC38A2 also acts as a selective target for inhibiting growth of Gln-dependent breast cancer cell lines (PMID: 33028955).

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



37°C incubated and boiled mouse brain tissues were subjected to SDS PAGE followed by western blot with 25928-1-AP (SNAT2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 25928-1-AP (SNAT2 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 25928-1-AP (SNAT2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).