For Research Use Only

Serpin F1/PEDF Polyclonal antibody

Catalog Number:26045-1-AP 3 Publications





Basic Information	Catalog Number: 26045-1-AP	GenBank Accession Number: BC013984	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	350 ug/ml	5176	WB 1:500-1:2000
	Source:	UNIPROT ID:	IHC 1:500-1:2000
	Rabbit	P36955	
	Isotype: IgG Immunogen Catalog Number:	Full Name: serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 Calculated MW:	
	AG23296		
		418 aa, 46 kDa	
		Observed MW: 46 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA	WB : A375 cells, human serum	
	Cited Applications: WB, IF	IHC : human liver tissue,	
	Species Specificity:		
	human		
	Cited Species: human		
	TE buffer pH 9.0; (*) Alterno retrieval may be performed buffer pH 6.0		
Background Information	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of	RPINF1), is a multifunctional secreted p inctionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI.	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a
	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of I neurotrophic factor involved in ne in individuals with osteogenesis i	nctionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom. imperfecta, type VI.	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a a cells. Mutations in this gene were fou
	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of neurotrophic factor involved in ne in individuals with osteogenesis i	Antionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI.	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a
	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of I neurotrophic factor involved in ne in individuals with osteogenesis i Author Xiaoyan Fu	Antionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI. Pubmed ID Journal	secreted protein that is widely express undant secretory products of adipocytes mass and INS resistance. PEDF is a a cells. Mutations in this gene were fou Application
	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of I neurotrophic factor involved in ne in individuals with osteogenesis i Author Xiaoyan Fu Hao Zhang	Actionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abo PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI. Pubmed ID Journal 31113934 Cell Death Dis	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a a cells. Mutations in this gene were fou Application WB
Notable Publications	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of neurotrophic factor involved in ne in individuals with osteogenesis i Author Xiaoyan Fu Hao Zhang Lele Li	Actionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI. Pubmed ID Journal 31113934 Cell Death Dis 34278806 Hum Gene Ther	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a a cells. Mutations in this gene were fou Application WB WB
Background Information Notable Publications	tumorigenic, and neurotrophic fur including by osteoblasts and oste and circulating concentrations of I neurotrophic factor involved in ne in individuals with osteogenesis i Author Xiaoyan Fu Hao Zhang	actionsis. It is an approximately 50-kDa oclasts. PEDF is also one of the most abu PEDF correlate positively with body fat euronal differentiation in retinoblastom imperfecta, type VI. <b>Pubmed ID</b> Journal 31113934 Cell Death Dis 34278806 Hum Gene Ther 37070131 J Cell Mol Med	secreted protein that is widely express undant secretory products of adipocyte mass and INS resistance. PEDF is a a cells. Mutations in this gene were fou Application WB WB

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





human serum were subjected to SDS PAGE followed by western blot with 26045-1-AP (PEDF antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. A375 cells were subjected to SDS PAGE followed by western blot with 26045-1-AP (PEDF antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 26045-1-AP (PEDF antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).