For Research Use Only

IGSF2 Polyclonal antibody

Catalog Number: 26047-1-AP

1 Publications



Basic Information

Catalog Number: 26047-1-AP

Size: 1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG23297

GenBank Accession Number:

BC136484 GeneID (NCBI): 9398 **UNIPROT ID:** Q93033

Full Name: immunoglobulin superfamily,

Observed MW: 115 kDa

member 2

Applications

Tested Applications:

FC, IF/ICC, IHC, WB, ELISA

Cited Applications:

Species Specificity: human, mouse **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HCT 116 cells, RAW 264.7 cells, Jurkat cells

Purification Method:

WB 1:1000-1:5000 IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human placenta tissue, IF: RAW 264.7 cells,

Background Information

IGSF2, also known as CD101, is a low abundance protein. Expression of CD101 is found on monocytes, granulocytes and dendritic cells. In addition expression on T lymphocytes is important for cell activation. It plays a role as inhibitor of T-cells proliferation induced by CD3 and inhibits expression of IL2RA on activated T-cells and secretion of IL2. This antibody specifically recognises the 115 kDa protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuyang Liu	35350788	Front Immunol	IF

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26047-1-AP (IGSF2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 26047-1-AP (IGSF2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 26047-1-AP (IGSF2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed RAW 264.7 cells using 26047-1-AP (IGSF2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 RAW 264.7 cells were stained with 0.2ug IGSF2 antibody (26047-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.