

For Research Use Only

# UFL1 Polyclonal antibody

Catalog Number: 26087-1-AP

Featured Product

11 Publications



## Basic Information

Catalog Number:

26087-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG23404

GenBank Accession Number:

BC036379

GeneID (NCBI):

23376

UNIPROT ID:

O94874

Full Name:

KIAA0776

Observed MW:

80 kDa; 100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:200-1:800

## Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse

Cited Species:

human, mouse, bovine

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : HEK-293T cells,

IHC : mouse liver tissue,

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Meiqian Kuang	31721015	Cell Stress Chaperones	WB,IHC,IF
Chengmin Li	31078114	Mol Immunol	WB
Meiqian Kuang	32655766	Oxid Med Cell Longev	WB,IF

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 26087-1-AP (UFL1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 26087-1-AP (UFL1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HEK-293T cells were subjected to SDS PAGE followed by western blot with 26087-1-AP (KIAA0776 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.