### For Research Use Only

# TNF-alpha Polyclonal antibody

Catalog Number:26162-1-AP 8 Publications



**Basic Information** 

Catalog Number: 26162-1-AP

NM-013693 GeneID (NCBI): 21926

Full Name:

GenBank Accession Number:

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

Concentration: 1000 μg/ml Source:

> tumor necrosis factor Observed MW: 17 kDa

Immunogen Catalog Number:

AG24089

Rabbit

Isotype:

**Applications** 

**Tested Applications:** 

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

mouse

**Cited Species:** 

mouse, rat

Positive Controls:

WB: RAW 264.7 cells,

## **Background Information**

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus  $functions\ through\ its\ receptors\ TNFRSF1A/TNFR1\ and\ TNFRSF1B/TNFBR.\ This\ cytokine\ is\ involved\ in\ the\ regulation$ of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. Mouse and human TNF-alpha share 79% amino acid sequence identity. Unlike human TNF-alpha, the mouse form is glycosylated. In mouse deficiency of this gene is associated with defects in response to bacterial infection, with defects in forming organized follicular dendritic cell networks and germinal centers, and with a lack of primary B cell follicles.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jie Liu	39765279	Cell Signal	WB
Jiaxuan Li	39627688	Cell Mol Biol Lett	IHC
Shishi Shen	39603278	Neurobiol Dis	


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Untreated, LPS and Brefeldin A treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 26162-1-AP (Tnf alpha antibody) at dilution of 1:500 and incubated at room temperature for 1.5 hours.