

For Research Use Only

Draxin Polyclonal antibody

Catalog Number: 26342-1-AP

2 Publications



Basic Information

Catalog Number:

26342-1-AP

Concentration:

300 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG23666

GenBank Accession Number:

BC021286

GeneID (NCBI):

374946

UNIPROT ID:

Q8NBI3

Full Name:

chromosome 1 open reading frame

187

Observed MW:

32 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human

Positive Controls:

WB : mouse cerebellum tissue, SH-SY5Y cells, rat cerebellum tissue

Background Information

C1orf187, also named Draxin(Dorsal repulsive axon guidance protein), is a repulsive axon guidance cue. It is widely expressed in the developing hippocampus, glioma, and all histological types of the lung cancer cell. Draxin was shown to bind a 22-amino-acid peptide of the EGF domain of Netrin-1, and through Draxin/Netrin-1 interaction, Draxin may act as a secreted Netrin-1 antagonist. C1orf187 can inhibit or repel neurite outgrowth from dorsal spinal cord. It inhibits the stabilization of cytosolic beta-catenin (CTNNB1) via its interaction with LRP6, thereby acting as an antagonist of Wnt signaling pathway (PMID: 29622850).

Notable Publications

Author	Pubmed ID	Journal	Application
Jia-hao Bao	36045691	Front Immunol	WB
Barbara Łasut-Szyska	38460933	Chem Biol Interact	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse cerebellum tissue were subjected to SDS PAGE followed by western blot with 26342-1-AP (C1orf187 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.