

For Research Use Only

Plakophilin 2 Polyclonal antibody

Catalog Number: 26479-1-AP

2 Publications



Basic Information

Catalog Number:

26479-1-AP

Concentration:

750 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG24155

GenBank Accession Number:

BC094762

GeneID (NCBI):

5318

UNIPROT ID:

Q99959

Full Name:

plakophilin 2

Calculated MW:

837 aa, 93 kDa

Observed MW:

90-100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:6000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:200-1:800

IF/ICC: 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : Caco-2 cells, HepG2 cells, rat heart tissue

IP : HepG2 cells,

IHC : human ovary cancer tissue,

IF/ICC : Caco-2 cells,

Background Information

Plakophilin 2 also known as PKP2, belongs to the beta-catenin family. Plakophilin proteins contain numerous armadillo repeats, localize to cell desmosomes and nuclei, and participate in linking cadherins to intermediate filaments in the cytoskeleton (PMID: 10828611). Mutations in PKP2 are associated with Arrhythmogenic right ventricular dysplasia, familial, 9 (ARVD9)(PMID: 15489853).

Notable Publications

Author	Pubmed ID	Journal	Application
Hui Xu	39117967	J Neurooncol	WB
Xuan Lai	38840221	J Nanobiotechnology	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26479-1-AP (Plakophilin 2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-Plakophilin 2 (IP:26479-1-AP, 4ug; Detection:26479-1-AP 1:4000) with HepG2 cells lysate 1480 ug.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 26479-1-AP (Plakophilin 2 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed Caco-2 cells using Plakophilin 2 antibody (26479-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).