For Research Use Only

OAT1 Polyclonal antibody

Catalog Number:26574-1-AP 13 Publications





Basic Information	Catalog Number: 26574-1-AP	GenBank Accession Number: BC033682	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	850 µg/ml	9356	WB 1:500-1:2000	
	Source: Rabbit	UNIPROT ID: 04U2R8	IHC 1:50-1:500	
	Isotype: IgG Immunogen Catalog Number: AG24234	Full Name: solute carrier family 22 (organic anion transporter), member 6		
		Calculated MW: 62 kDa Observed MW: 60-65 and 70-80 kDa		
Applications	Tested Applications:	Positive Controls:		
	IHC, WB,ELISA Cited Applications:	WB: mouse brain tissue,		
	WB, IF	IHC : mouse brain tissue,		
	Species Specificity: human, mouse			
	Cited Species: human, rat, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	OAT1, also named as SLC22A6, has two GXXXG motifs in its transmembrane which is associated with protein processing and oligomerization of other proteins. OAT1 is involved in the renal elimination of endogenous and exogenous organic anions. It has been reported that OAT1 plays a key role in clearing endogenous metabolites, toxins and drugs from blood. OAT1 has some isoforms and the calculated MW ranges from 55 kDa to 62 kDa. 2657 1-AP antibody detects the 60-65 kDa (native forms) and 70-80 kDa (mature forms) protein in SDS-PAGE. (PMID: 21340049, 23389457, 23196129)			
Background Information	processing and oligomerization o exogenous organic anions. It has toxins and drugs from blood. OAT 1-AP antibody detects the 60-65 k	f other proteins. OAT1 is involved in th been reported that OAT1 plays a key m 1 has some isoforms and the calculate	ne renal elimination of endogenous and ole in clearing endogenous metabolites, d MW ranges from 55 kDa to 62 kDa. 265	
	processing and oligomerization o exogenous organic anions. It has toxins and drugs from blood. OAT 1-AP antibody detects the 60-65 k 21340049, 23389457, 23196129)	f other proteins. OAT1 is involved in th been reported that OAT1 plays a key m 1 has some isoforms and the calculate	ne renal elimination of endogenous and ole in clearing endogenous metabolites, d MW ranges from 55 kDa to 62 kDa. 265 ure forms) protein in SDS-PAGE. (PMID:	
	processing and oligomerization o exogenous organic anions. It has toxins and drugs from blood. OAT 1-AP antibody detects the 60-65 k 21340049, 23389457, 23196129)	f other proteins. OAT1 is involved in th been reported that OAT1 plays a key m 1 has some isoforms and the calculate Da (native forms) and 70-80 kDa (mat	ne renal elimination of endogenous and ole in clearing endogenous metabolites, d MW ranges from 55 kDa to 62 kDa. 265	
	processing and oligomerization o exogenous organic anions. It has toxins and drugs from blood. OAT 1-AP antibody detects the 60-65 k 21340049, 23389457, 23196129) Author Tianshu Gao	f other proteins. OAT1 is involved in the been reported that OAT1 plays a key re 1 has some isoforms and the calculate Da (native forms) and 70-80 kDa (mat Pubmed ID Journal	ne renal elimination of endogenous and ole in clearing endogenous metabolites, d MW ranges from 55 kDa to 62 kDa. 265 ure forms) protein in SDS-PAGE. (PMID: Application	
Background Information Notable Publications	processing and oligomerization o exogenous organic anions. It has toxins and drugs from blood. OAT 1-AP antibody detects the 60-65 k 21340049, 23389457, 23196129) Author Tianshu Gao Kuniko Hunter	f other proteins. OAT1 is involved in the been reported that OAT1 plays a key m 1 has some isoforms and the calculate Da (native forms) and 70-80 kDa (mat Pubmed ID Journal 36120294 Front Pharmacol	he renal elimination of endogenous and ole in clearing endogenous metabolites, d MW ranges from 55 kDa to 62 kDa. 265 ure forms) protein in SDS-PAGE. (PMID: Application WB	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse brain tissue were subjected to SDS PAGE followed by western blot with 26574-1-AP (OAT1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 26574-1-AP (OAT1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).