

For Research Use Only

Bcl2 Polyclonal antibody

Catalog Number: 26593-1-AP

Featured Product

1147 Publications



Basic Information

Catalog Number:

26593-1-AP

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG24260

GenBank Accession Number:

NM-009741

GeneID (NCBI):

12043

UNIPROT ID:

P10417

Full Name:

B-cell leukemia/lymphoma 2

Calculated MW:

26 kDa

Observed MW:

26 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:3000

IHC: 1:200-1:800

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF, CoIP

Species Specificity:

mouse, rat

Cited Species:

mouse, rat, pig, rabbit, canine, chicken, bovine, hamster, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : NIH/3T3 cells, C2C12 cells, mouse brain tissue, mouse heart tissue, mouse testis tissue, mouse kidney tissue, mouse spleen tissue, rat spleen tissue, rat spleen tissue., rat kidney tissue, rat testis tissue, mouse colon tissue

IHC : mouse spleen tissue, mouse brain tissue

FC (Intra) : NIH/3T3 cells,

Background Information

Bcl2 is a member of the B cell lymphoma 2 protein family. Members of this family regulate cell death in multiple cell types and can have either proapoptotic or antiapoptotic activities. The protein encoded by this gene inhibits mitochondrial-mediated apoptosis. This protein is an integral outer mitochondrial membrane protein that functions as part of signaling pathway that controls mitochondrial permeability in response to apoptotic stimuli. This protein may also play a role in neuron cell survival and autophagy. Abnormal expression and chromosomal translocations of this gene are associated with cancer progression in numerous tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Feng Zhu	36180975	Phytother Res	WB
Qian Wang	32999044	Mol Cancer Ther	WB
Siqing Ma	34658905	Front Physiol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26593-1-AP (Bcl2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 26593-1-AP (Bcl2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 26593-1-AP (Bcl2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Bcl2 antibody (26593-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



1×10^6 NIH/3T3 cells were intracellularly stained with 0.25 μ g Bcl2 Polyclonal antibody (26593-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 μ g rabbit IgG isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.