For Research Use Only

APTX Polyclonal antibody

Catalog Number: 26747-1-AP 1 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC104881

Antigen affinity purification

26747-1-AP Size:

GeneID (NCBI):

Recommended Dilutions:

650 ug/ml Source:

54840

WB 1:500-1:2000 IHC 1:50-1:500

Purification Method:

Rabbit Isotype: **UNIPROT ID:** Q7Z2E3

Full Name:

aprataxin

Immunogen Catalog Number:

Calculated MW:

342 aa, 39 kDa

IF/ICC 1:50-1:500

AG24945

Observed MW: 39 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

WB: HEK-293T cells, HepG2 cells

Species Specificity:

IHC: human intrahepatic cholangiocarcinoma tissue,

human

human ovary cancer tissue

IF/ICC: HepG2 cells,

Positive Controls:

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

APTX, also named as Aprataxin, is a nuclear protein, present in both the nucleoplasm and the nucleolus, which is a member of the histidine triad (HIT) superfamily. APTX is involved in DNA single-strand break repair, mediating protein-protein interactions with molecules responding to DNA damage. APTX contains three conserved domains: an N-terminal forkhead-associated (FHA) domain which mediates protein-protein interactions, a HIT domain that is similar to Hint, and a C-terminal zinc finger domain. Loss of function mutations in APTX, the gene encoding for Aprataxin, destabilize the Aprataxin protein and result in a rare neurological disorder known as ataxia-oculomotor apraxia, characterized by abnormal movements of the head and eyes.

Notable Publications

Author	Pubmed ID	Journal	Application
Fan Meng	39368091	Cell Rep	

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using APTX antibody (26747-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



HEK-293T cells were subjected to SDS PAGE followed by western blot with 26747-1-AP (APTX Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 26747-1-AP (APTX antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 26747-1-AP (APTX antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).