For Research Use Only

CXCR3 Polyclonal antibody

Catalog Number: 26756-1-AP

Featured Product

16 Publications



Basic Information

Catalog Number: GenBank Accession Number: 26756-1-AP BC034403 GeneID (NCBI): Concentration: 750 ug/ml 2833 **UNIPROT ID:** Source: Rabbit P49682

chemokine (C-X-C motif) receptor 3

Calculated MW: Immunogen Catalog Number: AG25056 368 aa, 41 kDa Observed MW:

70 kDa

Full Name:

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC, IF

Isotype:

Species Specificity: human, mouse **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, Jurkat cells, mouse liver tissue

Purification Method:

WB 1:500-1:3000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: mouse kidney tissue,

Background Information

CXCR3 (C-X-C chemokine receptor type 3) is a G-protein-coupled seven-transmembrane domain chemokine receptor that is expressed on the surface of a number of cell types, including activated CD4+ and CD8+ T cells, NK and NK-T cells, plasmacytoid dendritic cells, and some B cells. CXCR3 is activated by three related chemokines (CXCL9, CXCL10, and CXCL11) and plays an important role in effector T-cell and NK cell trafficking. CXCR3 has three isoforms termed CXCR3-A, CXCR3-B, and CXCR3-alt with the calculated molecular mass of 41kDa, 46kDa, and 29kDa respectively. The apparent molecular mass of CXCR3 detected by some scientists is 70kDa, but 45kDa by others. (PMID: 16847335, 15150261, 22885102, 19151743)

Notable Publications

Author	Pubmed ID	Journal	Application
Yuejiao Xin	36078141	Cells	WB,IHC
Heng Li	30883697	Br J Pharmacol	IF
Wei-Chih Lien	36684074	Bioeng Transl Med	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 26756-1-AP (CXCR3 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 26756-1-AP (CXCR3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).