### For Research Use Only

# CDK12/CRKRS Polyclonal antibody

Catalog Number: 26816-1-AP

Featured Product

9 Publications



**Purification Method:** 

WB 1:2000-1:16000 IF/ICC 1:300-1:1200

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 26816-1-AP
 BC150265

 Concentration:
 GeneID (NCBI):

 1200 ug/ml
 51755

 Source:
 UNIPROT ID:

 Rabbit
 Q9NYV4

Cdc2-related kinase, arginine/serine-

Immunogen Catalog Number:

AG25323 Calculated MW:

1490 aa, 164 kDa Observed MW: 180-200 kDa

Full Name:

**Applications** 

Tested Applications: WB, IF/ICC, ELISA Cited Applications:

WB, IP

Isotype:

Species Specificity: human Cited Species: human, mouse **Positive Controls:** 

WB: HeLa cells, K-562 cells, Jurkat cells

IF/ICC: A431 cells,

## **Background Information**

CRKRS (Cdc2-related kinase, Arg/Ser), also named as cyclin-dependent kinase 12 (CKD12), belongs to CDC2/CDKX subfamily. CDK12 can form a heterodimeric complex with its activating partner, cyclin K, to regulate important cellular physiological processes that include transcriptional regulation, regulation of DNA damage repair genes such as BRCA1 and ATM, suppression of intronic polyadenylation, and susceptibility to PARP inhibitors (PMID: 30487607, 24240700, 31694772). CDK12 has 3 isoforms with the calculated molecular mass of 164/163/141 kDa, and the apparent molecular mass of 180-200 kDa due to some modifications (PMID 11683387, 22988298).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yan Liang	34686673	Cell Death Discov	WB
Hui Li	33429148	Eur J Cancer	WB
Mareike Polenkowski	36590164	iScience	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



WB result of CDK12/CRKRS antibody (26816-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CDK12/CRKRS transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 26816-1-AP (CDK12/CRKRS antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CDK12/CRKRS antibody (26816-1-AP) at dilution of 1:600 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).