

For Research Use Only

Cyclin D1 Polyclonal antibody

Catalog Number: 26939-1-AP

Featured Product

444 Publications



Basic Information

Catalog Number:

26939-1-AP

Concentration:

500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG25393

GenBank Accession Number:

BC000076

GeneID (NCBI):

595

UNIPROT ID:

P24385

Full Name:

cyclin D1

Calculated MW:

295 aa, 34 kDa

Observed MW:

34 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:750-1:3000

IF/ICC 1:400-1:1600

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig, canine, bovine, sheep, goat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : LNCaP cells, C6 cells, HeLa cells, MCF-7 cells, NIH/3T3 cells, Neuro-2a cells, mouse lung tissue

IHC : human ovary tumor tissue, human oesophagus cancer tissue, human lung cancer tissue, mouse spleen tissue, mouse pancreas tissue

IF/ICC : MCF-7 cells, NIH/3T3 cells

Background Information

CCND1 (cyclin D1), also known as PRAD1 or BCL1, belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. CCND1 forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. The CCND1 gene, located on 11q13 has been reported to be overexpressed in mantle cell lymphoma (MCL) due to the chromosomal translocation. CCND1 has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Over-expression of CCND1 is known to correlate with the early onset of cancer and risk of tumor progression and metastasis.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhou Yang	32993738	J Exp Clin Cancer Res	WB
Pengyun Li	36130940	Cell Death Dis	WB
Siwen Zhang	34546849	Bioengineered	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26939-1-AP (Cyclin D1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 26939-1-AP (Cyclin D1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Non-treated HeLa and thymidine treated HeLa cells were subjected to SDS PAGE followed by western blot with 26939-1-AP (Cyclin D1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blot with Alpha Tubulin antibody as loading control.




Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 26939-1-AP (Cyclin D1 antibody) at dilution of 1:1500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 26939-1-AP (Cyclin D1 antibody) at dilution of 1:1500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1X10⁶ SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human Cyclin D1 (26939-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP, Clone:) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Cyclin D1 antibody (26939-1-AP) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), (CL594-Phalloidin, red).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Cyclin D1 antibody (26939-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human Cyclin D1 (26939-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).