

For Research Use Only

F11R Polyclonal antibody

Catalog Number: 26941-1-AP



Basic Information

Catalog Number: 26941-1-AP	GenBank Accession Number: BC001533	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 50848	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9Y624	
Isotype: IgG	Full Name: F11 receptor	
Immunogen Catalog Number: AG25641	Calculated MW: 33 kDa	
	Observed MW: 33 kDa	

Applications

Tested Applications:

FC, IHC, WB, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HUVEC cells, human peripheral blood platelets, U-87 MG cells

IHC : human breast cancer tissue, human spleen tissue

Background Information

The F11 receptor (F11R) was first identified in human platelets as a 32/35 kDa protein duplex that serves as the receptor for a functional antibody that activates platelets. It seems to play a role in epithelial tight junction formation. It was also reported to function as a receptor for reovirus and ligand for the integrin LFA1.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 26941-1-AP (F11R antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 26941-1-AP (F11R antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ U-87 MG cells were surface stained with 0.8 ug Anti-Human F11R (26941-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.8 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were not fixed .



Various lysates were subjected to SDS PAGE followed by western blot with 26941-1-AP (F11R antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.