## For Research Use Only

## JAM-A/CD321 Polyclonal antibody

Catalog Number: 26941-1-AP



**Basic Information** 

Catalog Number:

26941-1-AP

BC001533

Concentration:

600 ug/ml

50848

Source:

UNIPROT ID:

Rabbit

Q9Y624

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

600 ug/ml

50848

VNIPROT ID:

Rabit

Q9Y624

Full Name:

F11 receptor

33 kDa
Observed MW:

33 kDa

Calculated MW:

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: human

AG25641

Immunogen Catalog Number:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification Recommended Dilutions:

Recommended Diluti WB 1:1000-1:4000 IHC 1:50-1:500

Positive Controls:

WB: HUVEC cells, human peripheral blood platelets,

U-87 MG cells

IHC: human breast cancer tissue, human spleen tissue

## **Background Information**

The F11 receptor (F11R) was first identified in human platelets as a 32/35 kDa protein duplex that serves as the receptor for a functional antibody that activates platelets. It seems to play a role in epithelial tight junction formation. It was also reported to function as a receptor for reovirus and ligand for the integrin LFA1.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.