### For Research Use Only

# Caspase 7 Polyclonal antibody

Catalog Number: 27155-1-AP

Featured Product

**60 Publications** 



**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number: 27155-1-AP BC015799 GeneID (NCBI): Concentration: 350 ug/ml 840 Source:

**UNIPROT ID:** Rabbit P55210 Full Name: Isotype: caspase 7, apoptosis-related cysteine

Immunogen Catalog Number:

AG25904

Calculated MW: 303 aa, 34 kDa

Observed MW: 35 kDa

peptidase

**Tested Applications:** WB, IF/ICC, IF-P, IP, ELISA

Cited Applications:

WB, IHC, IP Species Specificity: human, mouse

Cited Species: human, mouse, rat Positive Controls:

WB: Jurkat cells, HEK-293 cells, LNCaP cells, PC-3

**Purification Method:** 

WB 1:500-1:1000

protein lysate

IF-P 1:50-1:500

IF/ICC 1:200-1:800

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells, MCF-7 cells, mouse liver tissue

IP: HEK-293 cells, IF-P: mouse brain tissue, IF/ICC: MCF-7 cells,

## **Background Information**

Caspase 7(CASP7), like caspases 3 and 6, contains a short prodomain and, upon apoptotic induction, the 35 kDa  $proform\ is\ converted\ into\ a\ 32\ kDa\ intermediate\ or\ preactive\ form\ which\ is\ further\ processed\ into\ two\ active$ brain, which is up-regulated and activated after traumatic injury (PMID:15953353). Caspase-7 is classified as a member of the subgroup of cysteine proteases most related to the Caenorhabditis elegans factor CED-3, which also includes caspase-3, -6, and -9(PMID:9426061). The protein is involved in the activation cascade of caspases responsible for apoptosis execution.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lei Zhang	34592228	Life Sci	WB
Lei Zhang	36116140	Phytother Res	WB
Rongrong Liao	36139227	Animals (Basel)	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Untreated and Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 27155-1-AP (Caspase 7 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




WB result of Caspase 7 antibody (27155-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-Caspase 7 transfected HEK-




IP result of anti-Caspase 7 (IP:27155-1-AP, 4ug; Detection:27155-1-AP 1:1000) with HEK-293 cells lysate 2600 ug.




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Caspase 7 antibody (27155-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Caspase 7 antibody (27155-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Caspase 7 antibody (27155-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using Caspase 7 antibody (27155-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).