For Research Use Only

WNT7A Polyclonal antibody Catalog Number:27177-1-AP 1 Publications





Basic Information	Catalog Number: 27177-1-AP	er: GenBank Accession Number: BC008811		Purification Method: Antigen affinity purification
	Concentration:	GeneID (NCE	31):	Recommended Dilutions:
	550 ug/ml Source: Rabbit	7476 UNIPROT ID: O00755 Full Name: wingless-type MMTV integration sit family, member 7A		WB 1:1000-1:5000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IF/ICC 1:300-1:1200 site
	Isotype: IgG Immunogen Catalog Number: AG26060			
		Calculated MW: 39 kDa Observed MW: 30-39 kDa		
Applications	Tested Applications: WB, IF/ICC, IP, ELISA	Positive C		ontrols: 293 cells, mouse kidney tissue, rat kidney
	Cited Applications: tissue IF IP : mous Species Specificity:			2,5 certs, mouse money tissue, lot Mulley
			IP : mouse	kidney tissue,
			IF/ICC : HepG2 cells,	
Background Information	WNT7A belongs to the Wnt family. WNT7A plays an important role in embryonic development, including dorsal versus ventral patterning during limb development, skeleton development and urogenital tract development (PMI 16826533). WNT7A is required for central nervous system (CNS) angiogenesis and blood-brain barrier regulation (PMID:30026314). WNT7A is highly expressed in basal cells and is secreted in abundance by basal cells (PMID: 36318668).			
Notable Publications	Author	Pubmed ID	Journal	Application
	Yuan Gao	39402333	EMBO Rep	IF
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





IP result of anti-WNT7A (IP:27177-1-AP, 4ug; A Detection:27177-1-AP 1:2000) with mouse kidney tissue lysate 2160 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using WNT7A antibody (27177-1-AP) at dilution of 1:600 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).

Various lysates were subjected to SDS PAGE followed by western blot with 27177-1-AP (WNT7A antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.