For Research Use Only

BDH2 Polyclonal antibody

Catalog Number:27207-1-AP

Featured Product

3 Publications

Full Name:



Purification Method:

WB 1:1000-1:4000 IHC 1:100-1:400

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 27207-1-AP BC095414

Size: GeneID (NCBI): 56898

Source: UNIPROT ID: Rabbit Q9BUT1

3-hydroxybutyrate dehydrogenase,

Immunogen Catalog Number: type 2

AG24934 Observed MW: 27 kDa

Applications

Tested Applications: IHC, WB, ELISA
Cited Applications:

WB, IHC

Isotype:

Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: BxPC-3 cells, HeLa cells

IHC: human breast cancer tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaofei Gao	35499082	J Clin Invest	WB
Jia-Zhou Liu	32605589	J Exp Clin Cancer Res	IHC
Tianxiao Yang	29618368	J Hematol Oncol	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 27207-1-AP (BDH2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 27207-1-AP (BDH2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 27207-1-AP (BDH2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



WB result of BDH2 antibody (27207-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-BDH2 transfected HeLa cells.

Various lysates were subjected to SDS PAGE followed by western blot with 27207-1-AP (BDH2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.