### For Research Use Only

# RHOA Polyclonal antibody

Catalog Number: 27213-1-AP 1 Publications



**Purification Method:** 

WB 1:500-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: 27213-1-AP

Size:

BC005976 GeneID (NCBI):

GenBank Accession Number:

900 ug/ml 387
Source: UNIPROT ID:

Rabbit P61586 Isotype: Full Name:

IgG ras homolog gene family, member A

Immunogen Catalog Number: Calculated MW:

AG25873 22 kDa

Observed MW: 22 kDa

**Applications** 

Tested Applications: WB, IHC, ELISA

Cited Applications:

WB

Species Specificity: human, mouse Cited Species:

rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse kidney tissue,

IHC: human stomach tissue,

# **Background Information**

RhoA is a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. Overexpression of RhoA is associated with tumor cell proliferation and metastasis. RhoA signalling is critical to many cellular processes including migration, mechanotransduction, and is often disrupted in carcinogenesis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yang Yao	33760185	Mol Med Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 27213-1-AP (RHOA antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 27213-1-AP (RHOA antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).