For Research Use Only

KIF2A Polyclonal antibody

Catalog Number: 27218-1-AP

Featured Product

1 Publications

BC031828

3796

O00139 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:



Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 27218-1-AP

Size:
1100 µg/ml
Source:
Rabbit
Isotype:

IgG kinesin heavy chain member 2A

Immunogen Catalog Number:Calculated MW:AG25692679 aa, 77 kDaObserved MW:

80 kDa~100 kDa

Applications

Tested Applications: WB, IF/ICC, IHC, ELISA Cited Applications:

Species Specificity: human, mouse Cited Species:

IHC

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, HeLa cells

IHC: mouse brain tissue,

IF/ICC: A549 cells,

Background Information

Kinesin Family Member 2A (KIF2A), also known as Kinesin-2, is a member of the kinesin-13 superfamily protein. KIF2A is a microtubule-depolymerizing kinesin motor protein with essential roles in neural progenitor division and axonal pruning during brain development. KIF2A is involved in microtubule depolymerization at minus ends of the spindle to generate pole-migration forces for chromosome movement. KIF2A is a crucial factor in the invasion and metastasis of various tumors, such as gastric cancer(PMID: 32801897), hepatocellular carcinoma(PMID: 36277155).

Notable Publications

Author	Pubmed ID	Journal	Application
Hideko Hoshina	38447546	Pathobiology	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 27218-1-AP (KIF2A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of KIF2A antibody (27218-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-KIF2A transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed A549 cells using KIF2A antibody (27218-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Alpha Tubulin antibody (66031-1-Ig, Clone: 1E4C11, red).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 27218-1-AP (KIF2A antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).