

For Research Use Only

# KCTD10 Polyclonal antibody

Catalog Number: 27279-1-AP

Featured Product

4 Publications



## Basic Information

Catalog Number:

27279-1-AP

Size:

750 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG25971

GenBank Accession Number:

BC040062

GeneID (NCBI):

83892

UNIPROT ID:

Q9H3F6

Full Name:

potassium channel tetramerisation domain containing 10

Calculated MW:

313 aa, 35 kDa

Observed MW:

35 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

## Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, rat, mouse

Positive Controls:

WB: Jurkat cells, HEK-293 cells, HeLa cells, K-562 cells, MCF-7 cells, U2OS cells

## Background Information

KCTD10 is a member of the polymerase delta-interacting protein 1 (PDIP1) gene family, which consists of 3 members, PDIP1, KCTD10 and TNFAIP1. KCTD10 is inducible by TNF- $\alpha$ , interacts with PCNA and the small subunit (p50) of DNA polymerase  $\delta$ . Previous reports have shown that KCTD10 is essential for mammalian development. Further functional studies showed that KCTD10 interacts with proliferating cell nuclear antigen (PCNA) and plays an important role in DNA synthesis.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xin Wu	31511519	Nat Commun	WB
Yun Feng	36381427	Genet Res (Camb)	WB
Yue Liu	33553178	Front Cell Dev Biol	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 27279-1-AP (KCTD10 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.