

For Research Use Only

COX2/ Cyclooxygenase 2/ PTGS2 Polyclonal antibody

Catalog Number: 27308-1-AP

Featured Product

91 Publications



Basic Information

Catalog Number:

27308-1-AP

Concentration:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26174

GenBank Accession Number:

BC013734

GeneID (NCBI):

5743

UNIPROT ID:

P35354

Full Name:

prostaglandin-endoperoxide synthase
2 (prostaglandin G/H synthase and
cyclooxygenase)

Calculated MW:

604 aa, 68 kDa

Observed MW:

69 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:100-1:400

IF-P 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, canine

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : A549 cells, mouse lung tissue, THP-1 cells

IHC : human lung tissue, human spleen tissue

IF-P : mouse lung tissue,

Background Information

COX2 (Prostaglandin G/H synthase 2, PTGS2) mediates the formation of prostaglandins from arachidonate. Its subunit structure is homodimer. The fully N-glycosylated PTGS2 is 72-74 kDa and the aglycosylated is 66 kDa (PMID:19656660). It also expresses a band of 39 kDa after unspecific cleavage (PMID:17509125). The 50 kDa band of fragmented PTGS2 has also previously been detected in AD brains (PMID:14724276).

Notable Publications

Author	Pubmed ID	Journal	Application
Lin-Tao Xu	34601084	J Ethnopharmacol	WB
Koiti Aasumets	34622037	Biochem Biophys Rep	WB
Dewei He	36139502	Cells	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 27308-1-AP (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




A549 cells were subjected to SDS PAGE followed by western blot with 27308-1-AP (COX2/ Cyclooxygenase 2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung tissue slide using 27308-1-AP (COX2/ Cyclooxygenase 2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung tissue slide using 27308-1-AP (COX2/ Cyclooxygenase 2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse lung tissue using COX2/ Cyclooxygenase 2/ PTGS2 antibody (27308-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse lung tissue using COX2/ Cyclooxygenase 2/ PTGS2 antibody (27308-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).