For Research Use Only

Follistatin Polyclonal antibody Catalog Number: 27350-1-AP 1 Publications





Basic Information	Catalog Number: 27350-1-AP	GenBank Accession Number BC004107	Purification Method: Antigen affinity purification	
	Size: 1000 µg/ml	GenelD (NCBI): 10468	Recommended Dilutions: WB 1:500-1:1000	
	Source: Rabbit	UNIPROT ID: P19883	IF/ICC 1:200-1:800	
	lsotype: IgG	Full Name: follistatin		
	Immunogen Catalog Number: AG26401	Calculated MW: 33 aa, 7 kDa		
		Observed MW: 35 kDa		
Applications	Tested Applications: WB, IF/ICC, ELISA		tive Controls: A549 cells, Calu-3 cells, HeLa cells, U2OS cells CC : HeLa cells,	
	Cited Applications:			
	Species Specificity: human			
	Cited Species: mouse, rat			
Background Information	Follistatin (FST) is a member of the tissue growth factor β family and is a secreted glycoprotein that antagonizes many members of the family, including activin A, growth differentiation factor11, and myostatin. It binds activin A with high affinity and whose expression can be induced by activin A and several other pro-inflammatory cytokines Activin A-follistatin complexes are biologically inactive and bind to cell surface heparan sulphate-containing proteoglycans for internalisation and degradation. It has also been found to play a significant role in the management of skeletal muscle size and mass. It also has important roles in early embryonic development, differentiation of ovarian granulosa cells, liver fibrosis and polycystic ovarian syndrome.			
Notable Publications	Author	Pubmed ID Journal	Application	
	Ann Kuganathan	39300291 Hypertens R	les IHC	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Δ549 2150 kDa→ 1150 kDa→ 70 kDa→ 50 kDa→ 30 kDa→ 20 kDa→ 15 kDa→ 15 kDa→ 15 kDa→

Various lysates were subjected to SDS PAGE followed by western blot with 27350-1-AP (Follistatin antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Follistatin antibody (27350-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L). Various lysates were subjected to SDS PAGE followed by western blot with 27350-1-AP (Follistatin antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.