For Research Use Only

## TACSTD2/TROP2 Polyclonal antibody

Catalog Number: 27360-1-AP

2 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 27360-1-AP BC009409 GeneID (NCBI): Size: 750 μg/ml 4070 **UNIPROT ID:** Source: Rabbit P09758 Full Name: Isotype:

tumor-associated calcium signal

Immunogen Catalog Number: AG25769 Calculated MW: 36 kDa Observed MW:

45-55 kDa

transducer 2

**Applications** 

**Tested Applications:** IHC, WB, ELISA Cited Applications:

WB, IF

Species Specificity: human **Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HaCaT cells,

IHC: human breast cancer tissue, human prostate

**Purification Method:** 

WB 1:500-1:2400 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cancer tissue

## **Background Information**

Trophoblast cell surface antigen 2 (TROP2), also named as TACSTD2, is a transmembrane glycoprotein. TROP2 plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells, cornea and multi-stratified epithelia. TROP2 is overexpressed in most carcinomas and is involved in cancer proliferation, migration, invasion, and metastasis. Congenital mutations of TROP2 cause a rare autosomal recessive disease which may lead to blindness-the gelatinous drop-like corneal disease (PMID: 35688908).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Nadja Schöne	38453684	J Extracell Vesicles	WB
Wenjuan Tang	37271476	J Adv Res	WB,IF

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**



HaCaT cells were subjected to SDS PAGE followed by western blot with 27360-1-AP (TACSTD2 antibody) at dilution of 1:1200 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 27360-1-AP (TACSTD2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 27360-1-AP (TACSTD2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).