For Research Use Only

MUC2 Polyclonal antibody

Catalog Number: 27675-1-AP

56 Publications



Basic Information

Catalog Number: 27675-1-AP

Size: 500 µg/ml

Source: Rabbit M94132 GeneID (NCBI): 4583 Full Name:

mucin 2, oligomeric mucus/gel-

GenBank Accession Number:

Isotype: forming
IgG Calculated MW:
Immunogen Catalog Number: 540 kDa

AG25800

Applications

Tested Applications: IF/ICC,IF-P, IHC, ELISA Cited Applications: WB, IF, IHC

Species Specificity: Human, Mouse Cited Species:

human, rat, mosquito, mouse, pig, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method:

Antigen affinity purification Recommended Dilutions: IHC 1:1000-1:4000 IF 1:200-1:800

Positive Controls:

IHC : human colon tissue, mouse small intestine tissue

IF: HepG2 cells, human colon tissue, mouse colon tissue, rat small intestine tissue, SW480 cells

Background Information

This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. The protein polymerizes into a gel of which 80% is composed of oligosaccharide side chains by weight. Downregulation of this gene has been observed in patients with Crohn disease and ulcerative colitis.

Notable Publications

Author	Pubmed ID	Journal	Application
Nimra Zafar Siddiqui	36115959	Gut Pathog	IHC
Jianming Yang	36114582	Microbiome	IHC,IF
Shanshan Hu	34460244	J Agric Food Chem	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 27675-1-AP (MUC2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 27675-1-AP (MUC2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using MUC2 antibody (27675-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human colon tissue using MUC2 antibody (27675-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using MUC2 antibody (27675-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).