

For Research Use Only

MUC2 Polyclonal antibody, PBS Only

Catalog Number: 27675-1-PBS



Basic Information

Catalog Number:

27675-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG25800

GenBank Accession Number:

M94132

GeneID (NCBI):

4583

Full Name:

mucin 2, oligomeric mucus/gel-forming

Calculated MW:

540 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

IHC, IF/ICC, IF-P, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. The protein polymerizes into a gel of which 80% is composed of oligosaccharide side chains by weight. Downregulation of this gene has been observed in patients with Crohn disease and ulcerative colitis.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 27675-1-AP (MUC2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 27675-1-AP (MUC2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human colon tissue using MUC2 antibody (27675-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using MUC2 antibody (27675-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat small intestine tissue using MUC2 antibody (27675-1-AP) at dilution of 1:200 and Multi-rAb CoraLite® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004), smooth muscle actin specific antibody (67735-1-Ig, Clone: 1E9A11, green). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27675-1-PBS in a different storage



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using MUC2 antibody (27675-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.



1x10⁶ MCF-7 cells were intracellularly stained with 0.13 ug MUC2 Polyclonal antibody (27675-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.13 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 27675-1-PBS in a different storage buffer formulation.