### For Research Use Only

# SF3B1 Polyclonal antibody

Catalog Number: 27684-1-AP

**Featured Product** 

7 Publications



**Purification Method:** 

WB 1:1000-1:6000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number:

27684-1-AP

NM\_012433

Size:

GeneID (NCBI):

1000 µg/ml

23451

Source:

Rabbit

O75533

Isotype:

GeneID (NCBI):

Full Name:

splicing factor 3b, subunit 1, 155kDa

Immunogen Catalog Number: Calculated MW: 4G26689 146 kDa
Observed MW:

146-150 kDa

**Applications** 

Tested Applications: WB, IHC, ELISA Cited Applications:

WB

Species Specificity:

Human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: MCF-7 cells, HeLa cells, U2OS cells

IHC: human colon tissue,

# **Background Information**

SF3B1, also named as SAP155, is a 1304 amino acid protein, which belongs to the SF3B1 family. SF3B1 is a subunit of the splicing factor SF3B required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA. SF3B1 may also be involved in the assembly of the 'E' complex. SF3B1 belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Zixiang Wang	36271053	Nat Commun	WB
Yapu Li	33319461	EMBO Rep	WB
Tetsuro Ochi	36028755	Sci Rep	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




MCF-7 cells were subjected to SDS PAGE followed by western blot with 27684-1-AP (SF381 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 27684-1-AP (SF3B1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 27684-1-AP (SF3B1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 27684-1-AP (SF3B1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).