## For Research Use Only

# ABCG5 Polyclonal antibody

Catalog Number: 27722-1-AP 35 Publications



**Basic Information** 

Catalog Number:

27722-1-AP

Concentration:

850 µg/ml

Source:

Rabbit

Q9H222

Isotype:

GenBank Accession Number:

NM\_022436

GeneID (NCBI):

64240

UNIPROT ID:

Q9H222

Full Name:

ATP-binding cassette, sub-family G

AG26844 Calculated MW: 72 kDa
Observed MW:

68 kDa~72 kDa

(WHITE), member 5

**Applications** 

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC

Immunogen Catalog Number:

Species Specificity: Human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Caco-2 cells, HepG2 cells, mouse colon tissue, mouse liver tissue, rat liver tissue

Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human liver tissue,

## **Background Information**

### Notable Publications

Author	Pubmed ID	Journal	Application
Yunying Huang	34579033	Nutrients	WB
Feng Zhao	34803510	Int J Biol Sci	WB
Yunxia Yang	31679333	J Agric Food Chem	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 27722-1-AP (ABCG5 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 27722-1-AP (ABCG5 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 27722-1-AP (ABCG5 antibody) at dilution of 1:200 (under 40x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 27722-1-AP (ABCG5 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.