

TNC/Tenascin-C Polyclonal antibody

Catalog Number: 27789-1-AP

Basic Information

Catalog Number: 27789-1-AP	GenBank Accession Number: BC151843	Purification Method: Antigen affinity purification
Size: 450 µg/ml	GeneID (NCBI): 3371	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: P24821	IHC 1:500-1:2000
Isotype: IgG	Full Name: tenascin C	
Immunogen Catalog Number: AG27122	Calculated MW: 2201 aa, 241 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : U-87 MG cells,

IHC : human liver cancer tissue, human breast cancer tissue, human lung cancer tissue

Background Information

Tenascin-C (TNC) is a large hexameric extracellular matrix glycoprotein of the tenascin family (PMID: 21818551). It is a multimodular protein containing multiple epidermal growth factor (EGF)-like repeats and fibronectin type III (FN III) domains (PMID: 1719530). Tenascin-C is highly expressed during embryonic development, particularly in the developing central nervous system, around motile cells and at epithelial-mesenchymal interaction sites (PMID: 25738825). In adult tissues, the expression and the distribution of TNC are typically limited under normal physiological conditions. It is upregulated during injury, inflammation, regeneration, and cancer (PMID: 7694605; 25120494). Tenascin-C is a diverse protein that can produce different functions including regulating cell adhesion, migration and proliferation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



U-87 MG cells were subjected to SDS PAGE followed by western blot with 27789-1-AP (TNC antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27789-1-AP (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27789-1-AP (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).