For Research Use Only

eRF3a/GSPT1 Polyclonal antibody

Catalog Number:28130-1-AP

Featured Product 1 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 28130-1-AP	GenBank Accession Number: BC009503	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	500 ug/ml	2935	WB 1:1000-1:4000	
	Source:	UNIPROT ID:	IHC 1:500-1:2000 IF/ICC 1:200-1:800	
	Rabbit	P15170		
	Isotype: IgG	Full Name: G1 to S phase transition 1		
	Immunogen Catalog Number: AG27963	Calculated MW: 68 aa, 4 kDa		
		Observed MW: 80 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA Cited Applications:		WB : A549 cells, HepG2 cells, PC -3 cells, SKOV -3 cells Hela cells	
	WB Species Specificity:	IHC : hum tissue	an ovary cancer tissue, human stomach	
	human, mouse, rat	iuman, mouse, rat IF/ICC : He		
	Cited Species: human			
	TE buffer pH 9.0; (*) Alterno retrieval may be performed buffer pH 6.0			
Background Informatior	The eukaryotic Release Factor 3 (e termination. Eukaryotic release fa regulation of the cell cycle at the stimulator of eRF1 in translation t degradation machinery in the reoc transcription regulation. eRF3a, al in translation termination in respe ERF1. eRF3a/GSPT1 exists some is eRF3a/GSPT1 as a BIR3-binding p binding proteins from extracts of I	actor 3 (eRF3) has many functions in er G1 to S phase transition, and regulatir ermination. It was also reported to pla valing of ribosomes in successive cycl so known as GSPT1, is one subunit of ponse to the termination codons UAA, L soforms with MV 69 kDa and 56 kDa. I rotein-Using a GST-BIR3 fusion protein numan cells and mouse tissues, we pro- band confirmed that it is a processed	Jkaryotic cells, such as controlling the g protein synthesis as a GTP dependent by a key role as an initiator of the mRNA es of translation, and probably also in eRF3(PMID:15917414, 12923185). It involve IAG and UGA and stimulates the activity of	
Background Informatior	The eukaryotic Release Factor 3 (e termination. Eukaryotic release fa regulation of the cell cycle at the stimulator of eRF1 in translation t degradation machinery in the recy transcription regulation. eRF3a, al in translation termination in respe ERF1. eRF3a/GSPT1 exists some is eRF3a/GSPT1 exists some is eRF3a/GSPT1 as a BIR3-binding p binding proteins from extracts of I weights 23, 38 and 80 kDa. 80 kDa lacking the first 69 residues(PMID	actor 3 (eRF3) has many functions in er G1 to S phase transition, and regulatir ermination. It was also reported to pla valing of ribosomes in successive cycl so known as GSPT1, is one subunit of ponse to the termination codons UAA, L soforms with MV 69 kDa and 56 kDa. I rotein-Using a GST-BIR3 fusion protein numan cells and mouse tissues, we pro- band confirmed that it is a processed	g protein synthesis as a GTP dependent by a key role as an initiator of the mRNA es of translation, and probably also in eRF3(PMID:15917414, 12923185). It involve IAG and UGA and stimulates the activity of dentification of a processed form of n as an affinity reagent to purify new IAP eviously isolated 3 proteins of molecular	
	The eukaryotic Release Factor 3 (d termination. Eukaryotic release fa regulation of the cell cycle at the stimulator of eRF 1 in translation t degradation machinery in the recy transcription regulation. eRF3a, al in translation termination in resp ERF 1. eRF3a/GSPT1 exists some is eRF3a/GSPT1 as a BIR3-binding p binding proteins from extracts of I weights 23, 38 and 80 kDa. 80 kDa lacking the first 69 residues(PMID Author	actor 3 (eRF3) has many functions in er G1 to S phase transition, and regulatir ermination. It was also reported to pla valing of ribosomes in successive cycl so known as GSPT1, is one subunit of ponse to the termination codons UAA, L soforms with MV 69 kDa and 56 kDa. I rotein-Using a GST-BIR3 fusion proteir numan cells and mouse tissues, we pro b band confirmed that it is a processed : 12865429).	Jkaryotic cells, such as controlling the g protein synthesis as a GTP dependent ay a key role as an initiator of the mRNA es of translation, and probably also in eRF3(PMID:15917414, 12923185). It involve IAG and UGA and stimulates the activity of dentification of a processed form of as an affinity reagent to purify new IAP eviously isolated 3 proteins of molecular form of the human GSPT1/eRF3a protein,	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A549 cells were subjected to SDS PAGE followed by western blot with 28130-1-AP (eRF3a/CSPT1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using eRF3a/GSPT1 antibody (28130-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 28130-1-AP (eRF3a/GSPT1 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of eRF3a/GSPT1 antibody (28130-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-eRF3a/GSPT1 transfected HepG2 cells.