For Research Use Only

Caspase 14 Polyclonal antibody

Catalog Number: 28136-1-AP



Basic Information

Catalog Number: 28136-1-AP

Concentration:
800 ug/ml
Source:
Rabbit
Isotype:

IgG caspase 14, apoptosis-related cysteine peptidase

Immunogen Catalog Number: cysteine peptidase AG27242 Calculated MW: 242 aa, 28 kDa

Observed MW: 35 kDa

BC069541

23581

P31944 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000

Applications

Tested Applications: WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: mouse skin tissue, rat skin tissue

Background Information

Caspase 14 belongs to the caspase family of cysteinyl aspartate-specific proteinases that frequently play a central role in apoptosis. Caspase 14 exist as inactive proenzymes that undergo proteolytic processing to produce large and small subunits (p20 and p10) at Ile152/Lys153 residues, which is different to other caspases (Asp residue). Caspase 14 plays a role in protein maturation of filaggrin, and possibly in DNA repair. Overexpression of Caspase 14 has been detected in several of epithelial malignancies, indicating that Caspase 14 could play a role in carcinogenesis and disease progression of cancers. Caspase 14 can binds the apoptosis-inducing factor (AIF) indicating that caspase 14 may be an anti-apoptotic protein. In the LADC specimens, two bands can be detected with MW of 35 kDa and 37 kDa which represent Caspase 14 and the phosphorylated Caspase 14(pCasp-14) respectively. (PMID:21567094, 12200134, 10203698, 17515931, 16061209). This antibody can recognise all the forms of Caspase 14.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28136-1-AP (Caspase 14 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.