

For Research Use Only

CXCR5 Polyclonal antibody

Catalog Number: 28239-1-AP



Basic Information

Catalog Number: 28239-1-AP	GenBank Accession Number: NM_001716	Purification Method: Antigen affinity purification
Size: 350 µg/ml	GeneID (NCBI): 643	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	UNIPROT ID: P32302	
Isotype: IgG	Full Name: chemokine (C-X-C motif) receptor 5	
Immunogen Catalog Number: AG28271	Calculated MW: 42 kDa	
	Observed MW: 33 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Jurkat cells, K-562 cells, mouse spleen tissue, rat spleen tissue
Species Specificity: Human, mouse, rat	

Background Information

Initially discovered in 1992 as Burkitt's lymphoma receptor 1, CXCR5 (chemokine receptor 5, also known as CD185) is a seven-transmembrane G protein-coupled receptor (GPCR) protein (PMID: 32994482). CXCR5 is essential for naïve B-cell migration to and maturation in the lymph nodes and spleen, as well as for cluster of differentiation 4 T-cell (TFH and TCM) homing and interaction with B cells (PMID: 21471443). Moreover, CXCR5 is involved in the regulation of neural stem cells (NSC) and neuronal regeneration in the mouse brain, neuronal differentiation, neurogenesis, gliogenesis, and synaptogenesis (PMID: 37460877).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28239-1-AP (CXCR5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



WB result of CXCR5 antibody (28239-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CXCR5 transfected Jurkat cells.