For Research Use Only

MFN2 Polyclonal antibody

Catalog Number: 28341-1-AP



Basic Information

Catalog Number: 28341-1-AP

Size: GeneID (NCBI):

1000 μ g/ml 9927

Source: ENSEMBL Gene ID:
Rabbit ENSG00000116688
Isotype: UNIPROT ID:
IgG 095140
Immunogen Catalog Number: Full Name:
AG28128 mitofusin 2

Calculated MW: 757 aa, 86 kDa Observed MW: 86 kDa

GenBank Accession Number:

BC017061

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications: IHC, WB,ELISA Species Specificity:

Human, Mouse, Rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells, HepG2 cells, HeLa cells, mouse

brain tissue, rat brain tissue IHC: mouse heart tissue,

Background Information

MFN2, also named as CPRP1 and KIAA0214, belongs to the mitofusin family. It is an Essential transmembrane GTPase, which mediates mitochondrial fusion. MFN2 acts independently of the cytoskeleton. It therefore plays a central role in mitochondrial metabolism and may be associated with obesity and/or apoptosis processes. Overexpression of MFN2 induces the formation of mitochondrial networks. It plays an important role in the regulation of vascular smooth muscle cell proliferation. Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 2A2 (CMT2A2). Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 6 (CMT6). Ubiquitinated forms of Mfn2 (mono- and polyubiquitinated) are present during mitophagy.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 28341-1-AP (MFN2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 28341-1-AP (MFN2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 28341-1-AP (MFN2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 28341-1-AP (MFN2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).