For Research Use Only

PRKAR2B Polyclonal antibody

Catalog Number: 28351-1-AP 1 Publications



Basic Information

Catalog Number: 28351-1-AP Size:

650 ug/ml
Source:
Rabbit
Isotype:
IgG

Immunogen Catalog Number:

AG28787

AU20/0/

GenBank Accession Number:

BC075800 GeneID (NCBI): 5577 UNIPROT ID: P31323

protein kinase, cAMP-dependent, regulatory, type II, beta

Calculated MW: 418 aa, 46 kDa

Observed MW: 46-50 kDa

Full Name:

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: Y79 cells, human testis tissue, mouse brain tissue, mouse eye tissue, rat brain tissue, mouse testis tissue

IP: mouse brain tissue,
IHC: human stomach tissue,

IF/ICC: Neuro-2a cells, SH-SY5Y cells

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Enyi Zhu	38332898	J Inflamm Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28351-1-AP (PRKAR2B antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using PRKAR2B antibody (28351-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using PRKAR2B antibody (28351-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-PRKAR2B (IP:28351-1-AP, 4ug; Detection:28351-1-AP 1:1000) with mouse brain tissue lysate 1080 ug.




Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using PRKAR2B antibody (28351-1-AP) at dilution of 1:400 and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



1X10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human PR(AR2B (28351-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 28351-1-AP (PRKAR2B antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 28351-1-AP (PRKAR2B antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).