

For Research Use Only

UPF1 Polyclonal antibody

Catalog Number: 28384-1-AP

Featured Product



Basic Information

Catalog Number:

28384-1-AP

Size:

450 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG29057

GenBank Accession Number:

BC039817

GeneID (NCBI):

5976

UNIPROT ID:

Q92900

Full Name:

UPF1 regulator of nonsense transcripts homolog (yeast)

Calculated MW:

123 kDa

Observed MW:

123-130 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, HeLa cells, Raji cells, SH-SY5Y cells, SKOV-3 cells

IHC : human breast cancer tissue,

Background Information

Up-Frameshift Suppressor 1 Homolog (UPF1) is the central factor in nonsense-mediated mRNA decay (NMD) and is also directly involved in telomere homeostasis, DNA replication, histone mRNA degradation and staufen-mediated mRNA decay (PMID: 29382845). It is a potential modulator of MALAT1 and that UPF1/MALAT1 pathway could be a therapeutic target for gastric cancer (PMID: 28942451). The molecular mass of UPF1 is 123-130 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28384-1-AP (UPF1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of UPF1 antibody (28384-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-UPF1 transfected HEK-293 cells.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 28384-1-AP (UPF1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).