

For Research Use Only

COP1 Polyclonal antibody

Catalog Number: 28395-1-AP



Basic Information

Catalog Number: 28395-1-AP	GenBank Accession Number: BC039723	Purification Method: Antigen affinity purification
Source: Rabbit	GeneID (NCBI): 64326	Recommended Dilutions: WB: 1:1000-1:4000
Isotype: IgG	UNIPROT ID: Q8NHY2	IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Immunogen Catalog Number: AG28163	Full Name: ring finger and WD repeat domain 2	IHC: 1:50-1:500
	Calculated MW: 731 aa, 80 kDa	
	Observed MW: 80-90 kDa	

Applications

Tested Applications: WB, IHC, IP, ELISA	Positive Controls: WB : HCT 116 cells, IP : HCT 116 cells, HeLa cells IHC : mouse heart tissue,
Species Specificity: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

COP1 (constitutive photomorphogenic 1) is also named Ring finger and WD domain 2 (RFWD2), RING finger protein 200 (RNF20). COP1 is an E3 ubiquitin ligase that modifies specific target proteins, priming their degradation via the ubiquitin proteasome system (PMID: 27335464). COP1 (also known as RFWD2) is an important suppressor of pathogenic *c/EBP β* -dependent gene expression programs in microglia (PMID: 32795415). COP1 is a tumour suppressor that negatively regulates ETV1, ETV4 and ETV5. The ubiquitin ligase COP1 is a tumour suppressor whose downregulation promotes prostatic epithelial cell proliferation and tumorigenesis (PMID: 21572435). COP1 has been identified as a novel and critical negative regulator of the tumor suppressor protein p53, and it influences p53-dependent transcription, apoptosis, cell cycle arrest and DNA repair (PMID: 16931761, PMID: 15103385). Recently, COP1 has been found to be overexpressed in human hepatocellular carcinoma, breast cancer and ovarian adenocarcinomas (PMID: 15492238, PMID: 20959491).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HCT 116 cells lysates were subjected to SDS PAGE followed by western blot with 28395-1-AP (COP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-COP1 (IP:28395-1-AP, 4ug; Detection:28395-1-AP 1:1000) with HCT 116 cells lysate 1920 ug.



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 28395-1-AP (COP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).