For Research Use Only

TBK1 Polyclonal antibody

Catalog Number: 28397-1-AP

Featured Product

65 Publications



Basic Information

Catalog Number:

28397-1-AP

BC034950

Concentration:

800 ug/ml

Source:

Rabbit

Q9UHD2

Isotype:

GenBank Accession Number:

BC034950

GeneID (NCBI):

29110

UNIPROT ID:

Q9UHD2

Full Name:

IgG TANK-binding kinase 1
Immunogen Catalog Number: Observed MW:

AG28894 84 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IP Species Specificity:

human

Cited Species:

human, mouse, rat, pig, monkey, chicken, bovine,

hamster

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HT-1080 cells, HCT 116 cells, HeLa cells, HepG2

Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cells, U2OS cells

IHC: human stomach cancer tissue, human liver cancer

tissue

IF/ICC: MCF-7 cells,

Background Information

TBK1, also named as tumor necrosis factor (TNF) receptor-associated factor NF-kB activator (TANK)-binding kinase 1 (TBK1), NF-kB-activating kinase (NAK), T2K, is a multimeric kinase that modulates inflammation and autophagy. It is a ubiquitously expressed serine-threonine kinase belonging to the 'noncanonical IkB kinases' (IKKs) recognized for its critical role in regulating type I IFN production (PMID: 27211305). And TBK1 is an important player in yet another critical cellular function, autophagy. This antibody is specific for TBK1.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaohua He	36147630	Biomed Res Int	WB
Yinghua Zhao	34471099	Signal Transduct Target Ther	WB
Qingxia Gao	36314820	J Virol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28397-1-AP (TBK1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 28397-1-AP (TBK1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 28397-1-AP (TBK1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of TBK1 antibody (28397-1-AP; 1:4000; room temperature for 1.5 hours) with wild-type and TBK1 knockout HeLa cells.




Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 28397-1-AP (TBK1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 28397-1-AP (TBK1 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using TBK1 antibody (28397-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 28397-1-AP (TBK1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).