For Research Use Only

## MSTO1 Polyclonal antibody Catalog Number: 28400-1-AP Featured Product

Featured Product





Basic Information	Catalog Number: 28400-1-AP	GenBank Accession Number: BC002535	Purification Method: Antigen affinity purification
	Size: 600 ug/ml Source: Rabbit	GeneID (NCBI): 55154 UNIPROT ID: Q9BUK6	Recommended Dilutions: WB 1:500-1:2000 IHC 1:500-1:2000
	Immunogen Catalog Number: AG29067	Observed MW: 57-62 kDa	
	Applications	Tested Applications: WB, IHC, ELISA	Positive C
Species Specificity: human			cells, HepG2 cells an ovary cancer tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	MSTO1, also named as LST005 and SLTP005, belongs to the misato family. Human MSTO1 is a ubiquitously expressed, evolutionary conserved protein with homologous regions to GTPases. MSTO1 is a cytoplasmic protein that modulates mitochondrial dynamics by promoting mitochondrial fusion. MSTO1 has 7 isoforms produced by alternative splicing. (PMID: 33612823)		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	1 50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 28400-1-AP (MSTO 1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

WB result of MSTO1 antibody (28400-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MSTO1 transfected HepG2 cells.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 28400-1-AP (MSTO1 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).